

## Erratum

---

### Erythrocyte membrane composition in patients with primary hypercholesterolemia

A. Vayá<sup>a,\*</sup>, M. Martínez Triguero<sup>a</sup>, E. Réganon<sup>b</sup>, V. Vila<sup>b</sup>, V. Martínez Sales<sup>b</sup>, E. Solá<sup>c</sup> and A. Hernández Mijares<sup>c</sup>

<sup>a</sup> *Hemorheology and Thrombosis Unit, La Fe University, Valencia, Spain*

<sup>b</sup> *Research Centre, La Fe University, Valencia, Spain*

<sup>c</sup> *Dr. Peset University Hospital, Valencia, Spain*

[*Clinical Hemorheology and Microcirculation* **40**(4) (2008), 289–294]

Author A. Ricart was omitted from the list of authors. Complete list of authors and their affiliations should be:

A. Vayá<sup>a,\*</sup>, M. Martínez Triguero<sup>a</sup>, E. Réganon<sup>b</sup>, V. Vila<sup>b</sup>, V. Martínez Sales<sup>b</sup>, E. Solá<sup>c</sup>, A. Hernández Mijares<sup>c</sup> and A. Ricart<sup>d</sup>

<sup>a</sup> *Hemorheology and Thrombosis Unit, La Fe University, Valencia, Spain*

<sup>b</sup> *Research Centre, La Fe University, Valencia, Spain*

<sup>c</sup> *Dr. Peset University Hospital, Valencia, Spain*

<sup>d</sup> *Department of Applied Linguistics, Polytechnic University of Valencia, Valencia, Spain*

Additionally, the name of Dr. A. Ricart should be removed from the Acknowledgements.

Furthermore, 2 references were missing from the list of references and should be added:

[8] K. Michalska-Malecka, L. Slowinska, M. Dorecka and W. Romaniuk, Correlations in some pathogenetic factors and values of hemorheological parameters in age-related macular degeneration, *Clin. Hemorheol. Microcirc.* **38**(3) (2008), 209–216.

[9] M.I. Spengler, S.M. Bertoluzzo, G. Catalani and M.L. Rasia, Study on membrane fluidity and erythrocyte aggregation in equine, bovine and human species, *Clin. Hemorheol. Microcirc.* **38**(3) (2008), 171–176.

Consequently, all references in this list and in the text will move up 2 places from reference [10] onward.

The Introduction of the article should read:

### 1. Introduction

The influence of plasmatic lipids on rheological blood behaviour have been extensively investigated [1–7]. Different studies have found that patients with familial hypercholesterolemia have a higher cholesterol/phospholipid (Chol/Ph) ratio in their red cell membranes [8–11], which is possibly associated with a high plasma cholesterol and the corresponding increase in the lipoproteins that exchange LDL cholesterol with that of the red cell membrane [12].