

Erratum

Conservation of receptor antagonist anti-tumor activity by epidermal growth factor receptor antibody expressed in transgenic corn seed

Dale L. Ludwig, Larry Witte, Daniel J. Hicklin, Marie Prewett, Rajiv Bassi, Douglas Burtrum, Daniel S. Pereira, Xenia Jimenez, Floyd Fox, Babita Saxena, Qinwei Zhou, Yuemei Ma, Xiaoqiang Kang, Dipa Patel, Michael Barry, Paul Kussie, Zhenping Zhu, Douglas A. Russell, William L. Petersen, Thomas P. Jury, Fernando Gaitan-Gaitan, Daniel L. Moran, Xavier Delannay, Bradley S. Storrs, Jacob Tou, Mark E. Zupec, Karen S. Gustafson, John McIntyre, S. Joseph Tarnowski and Peter Bohlen

[*Human Antibodies* 13(3) (2004), 81–90]

When this article was originally published, the values for clearance (ml/hr/kg) for Glycosylated Corn C225 and Aglycosylated Corn C225 were inadvertently switched in Table 1.

The corrected table is provided below.

Link to the original article can be found at: <http://iospress.metapress.com/content/grdwdxhnv07u4hnm/?p=7a7032d996374bc7ae33adc04ae476ea&pi=3>.

Table 1
Pharmacokinetic results of acute dose injection of cetuximab and corn-derived antibodies

Antibody	Half life (Hours)	Clearance (ml/hr/kg)
Cetuximab*	85 +/− 15	0.44 +/− 0.04
Glycosylated	43 +/− 3	1.16 +/− 0.27
Aglycosylated	109 +/− 1	0.35 +/− 0.01

*Historical half-life 100hours.