

Supplementary Data

Plasma Transthyretin as a Candidate Marker for Alzheimer's Disease

Latha Velayudhan^{a,d*}, Richard Killick^a, Abdul Hye^a, Anna Kinsey^a, Andreas Güntert^a, Steven Lynham^a, Malcolm Ward^b, Rufina Leung^a, Anbarasu Lourdasamy^c, Alvina W.M. To^a, John Powell^a and Simon Lovestone^a


^a*Department of Old Age Psychiatry, NIHR Specialist Biomedical Research Centre for Mental Health at the South London and Maudsley NHS Foundation Trust and King's College London and the MRC Centre for Neurodegeneration, King's College London, Institute of Psychiatry, London, UK*

^b*Proteome Sciences plc, Surrey, UK*

^c*MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, London, UK*

^d*Department of Health Sciences, University of Leicester, Leicester, UK*

Accepted 5 September 2011



Supplementary Figure 1. Scatter plot showing the correlation between the values of plasma transthyretin (TTR) concentration obtained by the two methods (ELISA and immunoblotting).

*Correspondence to: Dr. Latha Velayudhan, P.O. Box 70, Department of Old Age Psychiatry, Institute of Psychiatry, De Crespigny Park, Denmark Hill, London SE5 8AF, UK. Tel.: +44 (0)2078480508; Fax: +44 (0)2078480632; E-mails: Latha.Velayudhan@kcl.ac.uk and lv24@le.ac.uk.