

# Plasma Glial Fibrillary Acidic Protein Is Associated with $^{18}\text{F}$ -SMBT-1 PET: Two Putative Astrocyte Reactivity Biomarkers for Alzheimer's Disease

Pratishtha Chatterjee<sup>a,b,1,\*</sup>, Vincent Doré<sup>c,d,1</sup>, Steve Pedrini<sup>b,k</sup>, Natasha Krishnadas<sup>d,e</sup>, Rohith Thota<sup>a,f</sup>, Pierrick Bourgeat<sup>g</sup>, Milos D. Ikonovic<sup>h,i</sup>, Stephanie R. Rainey-Smith<sup>b,j,k,l</sup>, Samantha C. Burnham<sup>g</sup>, Christopher Fowler<sup>e</sup>, Kevin Taddei<sup>b,k</sup>, Rachel Mulligan<sup>d</sup>, David Ames<sup>m,n</sup>, Colin L. Masters<sup>e</sup>, Jürgen Fripp<sup>c</sup>, Christopher C. Rowe<sup>d,o</sup>, Ralph N. Martins<sup>a,b,k,p,2</sup> and Victor L. Villemagne<sup>d,q,2</sup> for the AIBL Research Group<sup>r</sup>

<sup>a</sup>Macquarie Medical School, Macquarie University, North Ryde, NSW, Australia

<sup>b</sup>School of Medical and Health Sciences, Edith Cowan University, Joondalup, WA, Australia

<sup>c</sup>The Australian eHealth Research Centre, CSIRO, Brisbane, QLD, Australia

<sup>d</sup>Department of Molecular Imaging & Therapy, Austin Health, Heidelberg, VIC, Australia

<sup>e</sup>The Florey Institute of Neuroscience and Mental Health, University of Melbourne, Parkville, VIC, Australia

<sup>f</sup>School of Biomedical Sciences and Pharmacy, College of Health, Medicine and Wellbeing, University of Newcastle, Newcastle, NSW, Australia

<sup>g</sup>Health and Biosecurity Flagship, Australian eHealth Research Centre, Herston, QLD, Australia

<sup>h</sup>Department of Neurology, University of Pittsburgh, Pittsburgh, PA, USA

<sup>i</sup>Geriatric Research Education and Clinical Center, VA Pittsburgh Healthcare System, Pittsburgh, PA, USA

<sup>j</sup>Centre for Healthy Ageing, Health Futures Institute, Murdoch University, Murdoch, WA, Australia

<sup>k</sup>Australian Alzheimer's Research Foundation, Sarich Neuroscience Research Institute, Nedlands, WA, Australia

<sup>l</sup>School of Psychological Science, University of Western Australia, Crawley, WA, Australia

<sup>m</sup>National Ageing Research Institute, Parkville, VIC, Australia

<sup>n</sup>Academic Unit for Psychiatry of Old Age, University of Melbourne, Melbourne, VIC, Australia

<sup>o</sup>The Florey Department of Neuroscience and Mental Health, University of Melbourne, Parkville, VIC, Australia

<sup>p</sup>School of Psychiatry and Clinical Neurosciences, University of Western Australia, Crawley, WA, Australia

<sup>q</sup>Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, USA

<sup>r</sup><https://aibl.csiro.au/>

Accepted 13 January 2023

Pre-press 9 February 2023

<sup>1</sup>These authors contributed equally to this work.

<sup>2</sup>Senior authors.

\*Correspondence to: Dr Pratishtha Chatterjee, Macquarie Medical School, Macquarie University, 75 Talavera Road, North Ryde, NSW 2109, Australia. E-mail: pratishtha.chatterjee@mq.edu.au.

**Abstract.**

**Background:** Astrocyte reactivity is an early event along the Alzheimer's disease (AD) continuum. Plasma glial fibrillary acidic protein (GFAP), posited to reflect astrocyte reactivity, is elevated across the AD continuum from preclinical to dementia stages. Monoamine oxidase-B (MAO-B) is also elevated in reactive astrocytes observed using  $^{18}\text{F}$ -SMBT-1 PET in AD.

**Objective:** The objective of this study was to evaluate the association between the abovementioned astrocyte reactivity biomarkers.

**Methods:** Plasma GFAP and  $\text{A}\beta$  were measured using the Simoa<sup>®</sup> platform in participants who underwent brain  $^{18}\text{F}$ -SMBT-1 and  $\text{A}\beta$ -PET imaging, comprising 54 healthy control (13  $\text{A}\beta$ -PET+ and 41  $\text{A}\beta$ -PET-), 11 mild cognitively impaired (3  $\text{A}\beta$ -PET+ and 8  $\text{A}\beta$ -PET-) and 6 probable AD (5  $\text{A}\beta$ -PET+ and 1  $\text{A}\beta$ -PET-) individuals. Linear regressions were used to assess associations of interest.

**Results:** Plasma GFAP was associated with  $^{18}\text{F}$ -SMBT-1 signal in brain regions prone to early  $\text{A}\beta$  deposition in AD, such as the supramarginal gyrus (SG), posterior cingulate (PC), lateral temporal (LT) and lateral occipital cortex (LO). After adjusting for age, sex, *APOE*  $\epsilon 4$  genotype, and soluble  $\text{A}\beta$  (plasma  $\text{A}\beta_{42/40}$  ratio), plasma GFAP was associated with  $^{18}\text{F}$ -SMBT-1 signal in the SG, PC, LT, LO, and superior parietal cortex (SP). On adjusting for age, sex, *APOE*  $\epsilon 4$  genotype and insoluble  $\text{A}\beta$  ( $\text{A}\beta$ -PET), plasma GFAP was associated with  $^{18}\text{F}$ -SMBT-1 signal in the SG.

**Conclusion:** There is an association between plasma GFAP and regional  $^{18}\text{F}$ -SMBT-1 PET, and this association appears to be dependent on brain  $\text{A}\beta$  load.

Keywords: Astrocyte reactivity, biomarkers,  $^{18}\text{F}$ -SMBT-1, glial fibrillary acidic protein, monoamine oxidase B, positron emission tomography

**INTRODUCTION**

The current core biomarkers for Alzheimer's disease (AD) include abnormal brain amyloid- $\beta$  ( $\text{A}\beta$ ) and tau accumulation assessed via positron emission tomography (PET), and abnormal cerebrospinal fluid (CSF)  $\text{A}\beta_{42}/\text{A}\beta_{40}$  ratio and phosphorylated-tau181 (p-tau181) levels assessed via immunoassays. However, due to the costs, limitations with widespread availability, and relative invasiveness associated with these core biomarkers, research has focused on the development of high accuracy surrogate blood-based biomarkers such as plasma  $\text{A}\beta_{42}/\text{A}\beta_{40}$  ratio [1, 2], plasma phosphorylated tau isoforms (p-tau181, p-tau231, p-tau217) [3–5], as well as plasma glial fibrillary acidic protein (GFAP) [6–8].

GFAP, an astrocyte cytoskeletal protein, is upregulated in reactive astrocytes [9]. Reactive astrocytes surround  $\text{A}\beta$  plaques, and GFAP upregulation in brain tissue is associated with  $\text{A}\beta$  plaque density [10–12]. It has been suggested that reactive astrocytes release GFAP into blood via astrocyte end-feet encompassing brain capillaries [13]. Indeed, plasma GFAP levels are elevated in individuals with high  $\text{A}\beta$  in the brain ( $\text{A}\beta$ +) compared to those with low  $\text{A}\beta$  in the brain ( $\text{A}\beta$ -) [6, 7, 14, 15]; and have demonstrated high accuracy in differentiating  $\text{A}\beta$ + from  $\text{A}\beta$ - participants, as assessed by PET, along the AD continuum [12]. Furthermore, plasma GFAP has a higher accuracy than CSF GFAP in distinguish-

ing  $\text{A}\beta$ + from  $\text{A}\beta$ - participants [8] and can predict progression from the mild cognitive impaired stage (MCI; prodromal AD) to AD [15].

In addition to expressing high levels of GFAP, reactive astrocytes overexpress the enzyme monoamine oxidase B (MAO-B), and therefore MAO-B has been used as a brain imaging marker for reactive astrocytes [16, 17]. Evidence of astrocyte reactivity using MAO-B tracers such as  $^{11}\text{C}$ -L-deprenyl-D2 ( $^{11}\text{C}$ -DED) has been reported in prodromal AD [16, 17]. Recently, a novel  $^{18}\text{F}$  MAO-B PET tracer, (S)-(2-methylpyrid-5-yl)-6-[(3- $^{18}\text{F}$ -fluoro-2-hydroxy)propoxy] quinoline ( $^{18}\text{F}$ -SMBT-1) with reversible binding kinetics [18, 19] demonstrated higher binding in the brain of  $\text{A}\beta$ + AD compared with  $\text{A}\beta$ - healthy controls (HC) [20]. Further, significantly higher  $^{18}\text{F}$ -SMBT-1 binding has been observed in  $\text{A}\beta$ + HC compared with  $\text{A}\beta$ -HC in several brain regions associated with early  $\text{A}\beta$  accumulation, including superior parietal, lateral occipital, posterior cingulate, supramarginal gyrus, and lateral temporal cortex [20–22], suggesting that astrocyte reactivity assessed using  $^{18}\text{F}$ -SMBT-1 PET is associated with early  $\text{A}\beta$  accumulation from the preclinical stage of AD.

While both plasma GFAP and  $^{18}\text{F}$ -SMBT-1 PET are potential astrocyte markers, previous studies have not yet investigated whether an association exists between the two astrocyte markers. Further, since activation of astrocytes has been reported to be in response to soluble  $\text{A}\beta$  and insoluble  $\text{A}\beta$  [23–26],

whether soluble or insoluble A $\beta$  influences this association is also of interest. Additionally, the comparison of the association of both astrocyte markers with cognitive performance is of interest. To address this gap in the literature, the current study investigated whether plasma GFAP and  $^{18}\text{F}$ -SMBT-1 PET are associated, accounting for soluble A $\beta$  (plasma A $\beta_{1-42}$ /A $\beta_{1-40}$  ratio) and/or aggregated A $\beta$  (A $\beta$  PET). While different astrocyte markers may reflect different astrocyte states, we hypothesize that plasma GFAP will be associated with  $^{18}\text{F}$ -SMBT-1 PET and this association is A $\beta$  dependent. Furthermore, given that plasma GFAP is associated with cognitive performance [6, 27], and a preliminary study found an association between  $^{18}\text{F}$ -SMBT-1 PET and cognitive performance [20], we also hypothesize that both astrocyte markers, plasma GFAP and  $^{18}\text{F}$ -SMBT-1 PET, will be inversely associated with cognitive performance, comparatively.

## METHODS

### Participants

Non-smoking participants in the current study were selected from the Australian Imaging, Biomarker & Lifestyle Flagship Study of Ageing (AIBL) cohort. Participant exclusion criteria are described in detail elsewhere [28, 29]. Briefly, exclusion criteria comprised history of non-AD dementia, schizophrenia, bipolar disorder, significant current (but not past) depression, Parkinson's disease, cancer (other than basal cell skin carcinoma) within the last two years, symptomatic stroke, uncontrolled diabetes, or current regular alcohol use exceeding two standard drinks per day for women or four per day for men. Participants were classified as either HC, individuals with AD based on the NINCDS-ADRDA criteria [30], or MCI based on Winblad et al. and Petersen et al. criteria, representing a high-risk state for the development of AD [31, 32]. For the current study, plasma GFAP levels were measured in 71 participants from a subset of 78 AIBL participants that underwent A $\beta$  PET and  $^{18}\text{F}$ -SMBT-1 PET, based on plasma sample availability for the corresponding SMBT-1 timepoint. Out of these 71 participants, 59 participants underwent  $^{18}\text{F}$ -SMBT-1 imaging within 12 months of blood collection, while 12 participants had  $^{18}\text{F}$ -SMBT-1 imaging over 12 months from blood collection (Supplementary Table 1).

This study was approved by the ethics committees of St. Vincent's Health and Austin Health in Mel-

bourne, Australia and Hollywood Private Hospital, Australia and Edith Cowan University in Perth, Australia and Macquarie University in Sydney, Australia. All participants provided written informed consent before participation.

### Neuropsychological tests

Participants underwent a comprehensive battery of neuropsychological tests. However, for this study, the primary measure used to examine cognitive performance was the Mini-Mental State Examination (MMSE; scores range from 0 to 30, indicating severe impairment to no impairment) [33] and the Clinical Dementia Rating scale (CDR; scores range from 0 to 3, indicating no impairment to severe impairment) [34].

### Measurement of plasma GFAP, A $\beta_{1-40}$ , and A $\beta_{1-42}$

EDTA plasma samples were fractionated from overnight fasted blood. Plasma samples were stored at  $-80^{\circ}\text{C}$  prior to thawing for measurement of GFAP, A $\beta_{1-40}$ , and A $\beta_{1-42}$  concentrations using the single-molecule array (Simoa<sup>®</sup>) platform using the Neurology 4-Plex E kit (QTX-103670, Quanterix, Billerica, MA) wherein calibrators were run in duplicates and samples were run in singlicates. Quality control (QC) was achieved by assessing the levels of the positive controls included in the Simoa kits. The analytical lowest limit of quantification was 11.6 pg/ml for GFAP, 1.51 pg/ml for A $\beta_{1-42}$  and 4.08 pg/ml for A $\beta_{1-40}$ . The average %CV of the two quality controls were 1.68% and 1.46% for GFAP, 1.28% and 1.06% for A $\beta_{1-42}$  and 0.2% and 2.19% for A $\beta_{1-40}$ , respectively.

### Determination of Apolipoprotein E (APOE) $\epsilon 4$ carrier status

The APOE  $\epsilon 4$  genotype is a known genetic risk factor for AD. The APOE genotype was determined from purified genomic DNA extracted from 0.5 ml whole blood. Each sample was genotyped for the presence of the three APOE variants ( $\epsilon 2$ ,  $\epsilon 3$ , and  $\epsilon 4$ ) based on TaqMan SNP genotyping assays for rs7412 (C 904973) and rs429358 (C 3084793) as per the manufacturer's instructions (AB Applied Biosystems by Life Technologies, Scoresby, VIC, Australia). Five percent of the samples were genotyped in duplicate

and 100% inter- and intra-assay concordance was observed.

### PET imaging

PET studies were conducted at the Department of Molecular Imaging & Therapy, Austin Health in Melbourne, Australia. All participants underwent A $\beta$  PET imaging with either  $^{18}\text{F}$ -Flutemetamol (FLUTE,  $n=3$ ),  $^{18}\text{F}$ -Florbetapir (FBP,  $n=2$ ), or  $^{18}\text{F}$ -NAV4694 (NAV,  $n=66$ ) to determine neocortical A $\beta$  load. NAV and FBP were synthesized in-house, as previously reported [35–37]. FLUTE was manufactured by Cyclotek Pty Ltd (<http://www.cyclotek.com>). The NAV and FBP PET scan acquisition consisted of 20 min ( $4 \times 5$  min) dynamic scans acquired at 50 min after an intravenous bolus injection of 185 MBq ( $\pm 10\%$ ) of NAV or FBP. Similarly, the participants who received FLUTE also underwent a 20 min ( $4 \times 5$  min) PET acquisition starting at 90 min after injection of 185 MBq ( $\pm 10\%$ ) of FLUTE. All A $\beta$  imaging results were expressed in Centiloids (CL) [35, 36, 38, 39]. A $\beta$  PET scans were spatially normalized and surface projected using CapAIBL [39]. The standard CL method was applied to determine A $\beta$  burden [38]. A CL value  $>20$  was selected to determine a high A $\beta$  (A $\beta$ +) scan [40, 41].

$^{18}\text{F}$ -SMBT-1 was synthesized in-house in the Department of Molecular Imaging & Therapy, Austin Health, as previously described [19].  $^{18}\text{F}$ -SMBT-1 yielded a greater than 95% radiochemical purity after high performance liquid chromatography purification, with an average decay-corrected radiochemical yield of 40% and a molar activity at the end of  $^{18}\text{F}$ -SMBT-1 synthesis  $>400$  GBq/ $\mu\text{mol}$ .  $^{18}\text{F}$ -SMBT-1 PET studies were acquired as a 20-min emission scan ( $4 \times 5$  min frames) starting 60 min after an intravenous bolus injection of 185 ( $\pm 10\%$ ) MBq of  $^{18}\text{F}$ -SMBT-1.  $^{18}\text{F}$ -SMBT-1 PET images were also spatially normalized and surface projected using CapAIBL [42, 43], and regional standard uptake values (SUV) at 60–80 min post injection were normalized using the subcortical white matter (WM) as reference region to generate cortical and subcortical semiquantitative SUV ratios (SUVR). We selected five regions of interest (ROI): supramarginal gyrus, the posterior cingulate, the lateral temporal, the lateral occipital and the superior parietal that were significant in our previous report [20].

No correction for partial volume effects was applied to the PET data.

### Statistical analyses

ROI and vertex-wise linear regression analyses were carried out using plasma GFAP levels as the dependent variable and  $^{18}\text{F}$ -SMBT-1 SUVR, A $\beta$  PET CL, plasma A $\beta_{1-42}$ /A $\beta_{1-40}$  ratio, age, and sex as the independent variables to understand the association between plasma GFAP and  $^{18}\text{F}$ -SMBT-1 signal after adjusting for confounding variables in all participants ( $N=71$ ). Similar analyses were carried out in participant subsets that: underwent  $^{18}\text{F}$ -SMBT-1 imaging within 12 months of blood collection ( $N=59$ ), were within the AD continuum (HC A $\beta$ +, MCI A $\beta$ +, AD A $\beta$ +,  $N=21$ ) or that comprised cognitively impaired participants ( $N=17$ ). Linear regression analyses were also carried out using A $\beta$  PET CL as the dependent variable and plasma GFAP,  $^{18}\text{F}$ -SMBT-1 SUVR, plasma A $\beta_{1-42}$ /A $\beta_{1-40}$  ratio, age, and sex as the independent variables to understand the association between plasma GFAP and A $\beta$  PET CL after adjusting for confounding variables. Years of education was not included in the above statistical models, because this information was not available for all study participants. Associations for MMSE with age and A $\beta$  PET CL adjusted plasma GFAP and  $^{18}\text{F}$ -SMBT-1 SUVR were carried out using Spearman's correlation ( $\rho$ ). Shapiro-Wilk tests were used to check normal distributions of model residuals. All analyses were carried out using IBM<sup>®</sup> SPSS<sup>®</sup> (v27).  $p < 0.05$  was considered significant.

## RESULTS

Table 1 shows participant characteristics including age, sex, and *APOE*  $\epsilon 4$  carrier status, MMSE scores, brain A $\beta$  PET load and plasma levels of GFAP and A $\beta_{1-42}$ /A $\beta_{1-40}$  ratio. This study included 54 HC (13 A $\beta$ +, and 41 A $\beta$ -), 11 MCI (3 A $\beta$ +, and 8 A $\beta$ -), and 6 probable AD (5 A $\beta$ +, and 1 A $\beta$ -) participants.

Plasma GFAP was significantly associated with  $^{18}\text{F}$ -SMBT-1 PET signal in four of the five brain regions examined, including the supramarginal gyrus (Standardized beta;  $\beta=0.361$ ,  $p=0.002$ ), posterior cingulate ( $\beta=0.308$ ,  $p=0.009$ ), lateral temporal ( $\beta=0.299$ ,  $p=0.011$ ), and lateral occipital ( $\beta=0.313$ ,  $p=0.008$ ) regions before adjusting for any covariates (Table 2). After adjusting for age, sex, *APOE*  $\epsilon 4$  carrier status, and soluble A $\beta$ , plasma GFAP was significantly associated with  $^{18}\text{F}$ -SMBT-1 PET signal in all regions: supramarginal gyrus ( $\beta=0.323$ ,  $p=0.002$ ), posterior cingulate ( $\beta=0.262$ ,

Table 1  
Participant characteristics

	HC Aβ <sup>-</sup>	HC Aβ <sup>+</sup>	MCI Aβ <sup>-</sup>	MCI Aβ <sup>+</sup>	AD Aβ <sup>-</sup>	AD Aβ <sup>+</sup>
N (M/F)	41 (16/25)	13 (6/7)	8 (5/3)	3 (3/0)	1 (1/0)	5 (2/3)
Age y, mean ± SD (range)	76.12 ± 4.94 (63–86)	78.89 ± 5.52 (71–89)	71.83 ± 6.37 (58–80)	75.10 ± 2.76 (72–77)	70.00	77.98 ± 3.13 (75–83)
<i>APOE</i> ε4 carrier status (N (%))	12 (29.27)	9 (69.23)	2 (25)	1 (33.33)	0 (0)	4 (80)
MMSE (mean ± SD)	28.51 ± 1.52	28.62 ± 1.61	28.25 ± 1.67	26.33 ± 0.58	27.00	24.00 ± 5.00
CDR (mean ± SD)	0.05 ± 0.15	0.00 ± 0.00	0.31 ± 0.26	0.50 ± 0.00	0.50	0.70 ± 0.27
Plasma GFAP (pg/mL, mean ± SD)	121.61 ± 43.63	180.74 ± 53.62	97.91 ± 26.19	122.74 ± 80.89	75.44	297.10 ± 139.81
Plasma Aβ <sub>1-42</sub> /Aβ <sub>1-40</sub> ratio (mean ± SD)	0.061 ± 0.013	0.054 ± 0.009	0.061 ± 0.012	0.055 ± 0.010	0.052	0.045 ± 0.013
Aβ PET (centiloid, mean ± SD)	0.55 ± 8.0	57.54 ± 47.78	1.55 ± 6.47	75.27 ± 63.05	-3.80	122.54 ± 42.28
SMBT-1 PET SG (SUVR, mean ± SD)	1.17 ± 0.10	1.31 ± 0.14	1.19 ± 0.07	1.30 ± 0.06	1.07	1.34 ± 0.07
SMBT-1 PET PC (SUVR, mean ± SD)	1.24 ± 0.12	1.36 ± 0.19	1.26 ± 0.04	1.36 ± 0.16	1.18	1.42 ± 0.12
SMBT-1 PET LT (SUVR, mean ± SD)	1.23 ± 0.13	1.36 ± 0.12	1.24 ± 0.07	1.28 ± 0.03	1.16	1.33 ± 0.08
SMBT-1 PET LO (SUVR, mean ± SD)	0.97 ± 0.10	1.07 ± 0.12	0.95 ± 0.04	1.06 ± 0.04	0.89	1.20 ± 0.08
SMBT-1 PET SP (SUVR, mean ± SD)	0.96 ± 0.12	1.06 ± 0.14	0.98 ± 0.12	1.04 ± 0.13	0.93	1.09 ± 0.19
Months between blood collection and SMBT-1 PET (Mean (minimum, maximum))	3.57 (-5, 39)	4.25 (-4, 24)	0.79 (0, 3)	5.08 (-1, 17)	9	12.95 (1, 24)

Sex, age, Apolipoprotein E (*APOE*) ε4 allele status, Mini-Mental State Examination (MMSE) scores, plasma glial fibrillary acidic protein (GFAP) levels, plasma Aβ<sub>1-42</sub>/Aβ<sub>1-40</sub> ratios, brain Aβ PET and brain monoamine oxidase B expression assessed using SMBT-1 PET in the supramarginal gyrus (SG), posterior cingulate (PC), lateral temporal (LT), lateral occipital (LO), and superior parietal (SP) are presented in participants stratified by clinical status and Aβ PET<sup>-/+</sup> status. HC, healthy control; MCI, mild cognitively impaired; AD, Alzheimer's disease; PET, positron emission tomography, <sup>18</sup>F-SMBT-1, (S)-(2-methylpyrid-5-yl)-6-[(3-<sup>18</sup>F-fluoro-2-hydroxy)propoxy]quinoline; SUVR, standardized uptake value ratio. Plasma GFAP levels and SMBT-1 PET SG, PC, LT, LO, and SP SUVRs were higher in Aβ<sup>+</sup> participants (*n* = 21) compared with Aβ<sup>-</sup> participants (*n* = 50) after adjusting for age and sex (*p* ≤ 0.01). Out of these 71 participants, 59 participants underwent <sup>18</sup>F-SMBT-1 imaging within 12 months of blood collection, while 12 participants had <sup>18</sup>F-SMBT-1 imaging over 12 months from blood collection (Supplementary Table 1).

Table 2  
Association of plasma GFAP with <sup>18</sup>F-SMBT-1 PET

	Regional SMBT-1 PET β (p)	Plasma Aβ <sub>1-42</sub> /Aβ <sub>1-40</sub> β (p)	Aβ PET β (p)	Age β (p)	Sex β (p)	APOE ε4 carrier status β (p)
A. No covariates included						
Supramarginal gyrus	0.361 ( <b>0.002</b> )	-	-	-	-	-
Posterior cingulate	0.308 ( <b>0.009</b> )	-	-	-	-	-
Lateral temporal	0.299 ( <b>0.011</b> )	-	-	-	-	-
Lateral occipital	0.313 ( <b>0.008</b> )	-	-	-	-	-
Superior parietal	0.197 (0.099)	-	-	-	-	-
B. Inclusion of age, sex and APOE ε4 carrier status						
Supramarginal gyrus	0.331 ( <b>0.002</b> )	-	-	0.256 ( <b>0.010</b> )	0.408 (< <b>0.001</b> )	0.100 (0.341)
Posterior cingulate	0.264 ( <b>0.013</b> )	-	-	0.270 ( <b>0.008</b> )	0.387 (< <b>0.001</b> )	0.129 (0.226)
Lateral temporal	0.229 ( <b>0.029</b> )	-	-	0.281 ( <b>0.006</b> )	0.363 (< <b>0.001</b> )	0.145 (0.175)
Lateral occipital	0.271 ( <b>0.017</b> )	-	-	0.233 ( <b>0.026</b> )	0.410 (< <b>0.001</b> )	0.115 (0.292)
Superior parietal	0.200 (0.057)	-	-	0.271 ( <b>0.010</b> )	0.386 (< <b>0.001</b> )	0.182 (0.084)
C. Inclusion of soluble Aβ, age, sex and APOE ε4 carrier status						
Supramarginal gyrus	0.323 ( <b>0.002</b> )	-0.245 ( <b>0.014</b> )	-	0.215 ( <b>0.026</b> )	0.421 (< <b>0.001</b> )	0.038 (0.715)
Posterior cingulate	0.262 ( <b>0.011</b> )	-0.253 ( <b>0.013</b> )	-	0.227 ( <b>0.022</b> )	0.402 (< <b>0.001</b> )	0.063 (0.551)
Lateral temporal	0.238 ( <b>0.019</b> )	-0.264 ( <b>0.011</b> )	-	0.235 ( <b>0.018</b> )	0.379 (< <b>0.001</b> )	0.073 (0.488)
Lateral occipital	0.282 ( <b>0.010</b> )	-0.266 ( <b>0.009</b> )	-	0.185 (0.068)	0.428 (< <b>0.001</b> )	0.041 (0.701)
Superior parietal	0.229 ( <b>0.024</b> )	-0.281 ( <b>0.007</b> )	-	0.219 ( <b>0.030</b> )	0.409 (< <b>0.001</b> )	0.105 (0.310)
D. Inclusion of insoluble Aβ, age, sex and APOE ε4 carrier status						
Supramarginal gyrus	0.215 ( <b>0.038</b> )	-	0.365 ( <b>0.001</b> )	0.195 ( <b>0.038</b> )	0.425 (< <b>0.001</b> )	-0.023 (0.821)
Posterior cingulate	0.190 (0.052)	-	0.402 (< <b>0.001</b> )	0.194 ( <b>0.039</b> )	0.418 (< <b>0.001</b> )	-0.026 (0.802)
Lateral temporal	0.174 (0.068)	-	0.415 (< <b>0.001</b> )	0.199 ( <b>0.036</b> )	0.402 (< <b>0.001</b> )	-0.023 (0.829)
Lateral occipital	0.142 (0.192)	-	0.395 (< <b>0.001</b> )	0.183 (0.060)	0.421 (< <b>0.001</b> )	-0.015 (0.885)
Superior parietal	0.149 (0.118)	-	0.421 (< <b>0.001</b> )	0.191 ( <b>0.046</b> )	0.420 (< <b>0.001</b> )	0.004 (0.972)
E. Inclusion of soluble Aβ, insoluble Aβ, age, sex and APOE ε4 carrier status						
Supramarginal gyrus	0.226 ( <b>0.027</b> )	-0.171 (0.079)	0.310 ( <b>0.007</b> )	0.176 (0.058)	0.431 (< <b>0.001</b> )	-0.048 (0.640)
Posterior cingulate	0.198 ( <b>0.041</b> )	-0.167 (0.088)	0.351 ( <b>0.002</b> )	0.176 (0.060)	0.424 (< <b>0.001</b> )	-0.050 (0.630)
Lateral temporal	0.187 ( <b>0.048</b> )	-0.172 (0.080)	0.362 ( <b>0.001</b> )	0.179 (0.055)	0.408 (< <b>0.001</b> )	-0.408 (0.644)
Lateral occipital	0.170 (0.118)	-0.179 (0.074)	0.332 ( <b>0.005</b> )	0.158 (1.00)	0.432 (< <b>0.001</b> )	-0.044 (0.675)
Superior parietal	0.175 (0.065)	-0.185 (0.064)	0.362 ( <b>0.001</b> )	0.167 (0.076)	0.430 (< <b>0.001</b> )	-0.022 (0.828)

Linear regressions were used to perform analyses in all study participants (N=71). Age, sex, APOE ε4 carrier status, soluble Aβ (plasma Aβ<sub>1-42</sub>/Aβ<sub>1-40</sub> ratio), and insoluble Aβ (Aβ PET) were also added to the model as covariates. Natural log plasma GFAP values were used in the model to satisfy Shapiro-Wilk test of normality of the model residuals. 'β' represents the standardized coefficients and 'p' represents significance, with p<0.05 considered significant (in bold). GFAP, glial fibrillary acidic protein; PET, positron emission tomography; <sup>18</sup>F-SMBT-1, (S)-(2-methylpyridin-5-yl)-6-[(3-<sup>18</sup>F-fluoro-2-hydroxy)propoxy] quinoline.




Fig. 1A. Vertex-wise analyses for the association between (a.) plasma GFAP and  $^{18}\text{F}$ -SMBT-1 PET SUVR, (b.) plasma GFAP and insoluble A $\beta$  (A $\beta$  PET) and (c.)  $^{18}\text{F}$ -SMBT-1 PET SUVR and insoluble A $\beta$  (A $\beta$  PET). The color scale represents  $p$  values.




Fig. 1B. Graphical representation of correlations between (a.) plasma GFAP and  $^{18}\text{F}$ -SMBT-1 PET SUVR, (b.) A $\beta$  PET and plasma GFAP, and (c.) A $\beta$  PET and  $^{18}\text{F}$ -SMBT-1 PET SUVR.

$p=0.011$ ), lateral temporal ( $\beta=0.238, p=0.019$ ), lateral occipital ( $\beta=0.282, p=0.010$ ), and superior parietal ( $\beta=0.229, p=0.024$ ) regions. However, after adjusting for age, sex, *APOE*  $\epsilon 4$  carrier status, and insoluble A $\beta$ , plasma GFAP was significantly associated with  $^{18}\text{F}$ -SMBT-1 PET signal only in the supramarginal gyrus ( $\beta=0.215, p=0.038$ ), with a trend towards significance observed in the posterior cingulate ( $\beta=0.190, p=0.052$ ) and lateral temporal ( $\beta=0.174, p=0.068$ ) regions (Table 2). On adjusting for soluble A $\beta$ , insoluble A $\beta$ , age, sex, and *APOE*  $\epsilon 4$  carrier status, plasma GFAP was significantly associated with  $^{18}\text{F}$ -SMBT-1 PET signal in the supramarginal gyrus ( $\beta=0.226, p=0.027$ ), posterior cingulate ( $\beta=0.198, p=0.041$ ), and lateral temporal ( $\beta=0.187, p=0.048$ ), with a trend towards significance observed in the superior parietal ( $\beta=0.175, p=0.065$ ) region (Fig. 1, Table 2). Similar observations from analyses in participants that underwent

$^{18}\text{F}$ -SMBT-1 imaging within 12 months of blood collection ( $N=59$ ) are presented in Supplementary Table 2.

No significant association was observed between plasma GFAP and SMBT-1 PET in participants within the AD continuum ( $N=21$ , Supplementary Table 3A). In cognitively impaired participants ( $N=17$ ), a significant (or trend towards significant) association was found (Supplementary Table 3B). When limiting the analysis to the healthy control group, no significant association was observed between plasma GFAP and  $^{18}\text{F}$ -SMBT-1 PET before and after adjusting for soluble A $\beta$ , insoluble A $\beta$ , age, sex, and *APOE*  $\epsilon 4$  carrier status.

Further, as expected, insoluble A $\beta$  PET was significantly associated with plasma GFAP, soluble A $\beta$  and  $^{18}\text{F}$ -SMBT-1 PET signal, after adjusting for confounding variables (Table 3 and Supplementary Table 4).

Table 3  
Association between insoluble A $\beta$  (measured via PET) with plasma GFAP

Plasma GFAP $\beta$ ( <i>p</i> )	Regional SMBT-1 PET $\beta$ ( <i>p</i> )	Plasma A $\beta_{1-42}$ /A $\beta_{1-40}$ $\beta$ ( <i>p</i> )	Age $\beta$ ( <i>p</i> )	Sex $\beta$ ( <i>p</i> )	APOE $\epsilon 4$ carrier status $\beta$ ( <i>p</i> )
0.417 (<0.001)	0.346 ( <b>0.001</b> )	-0.262 ( <b>0.010</b> )	0 (0.999)	-0.259 ( <b>0.018</b> )	0.081 (0.419)
0.446 (<0.001)	0.294 ( <b>0.006</b> )	-0.260 ( <b>0.013</b> )	-0.004 (0.966)	-0.288 ( <b>0.010</b> )	0.088 (0.392)
0.467 (<0.001)	0.260 ( <b>0.014</b> )	-0.269 ( <b>0.011</b> )	0.018 (0.858)	-0.316 ( <b>0.005</b> )	0.083 (0.430)
0.449 (<0.001)	0.329 ( <b>0.002</b> )	-0.287 ( <b>0.006</b> )	-0.044 (0.651)	-0.268 ( <b>0.016</b> )	0.078 (0.446)
0.439 (<0.001)	0.288 ( <b>0.006</b> )	-0.279 ( <b>0.008</b> )	-0.006 (0.953)	-0.269 ( <b>0.018</b> )	0.132 (0.193)

Linear regressions were used to perform the analyses in study participants (N=62) after removing outliers and utilizing natural log plasma GFAP values in the model to satisfy Shapiro-Wilk test of normality of model residuals.  $^{18}\text{F-SMBT-1}$  PET signal, soluble A $\beta$  (plasma A $\beta_{1-42}$ /A $\beta_{1-40}$ ), age, sex and APOE  $\epsilon 4$  carrier status were added to the model as covariates. Participants with standardized model residuals larger than an absolute value of  $\pm 2$  were considered as outliers. Removal of these outliers enabled normal distribution of model residuals. ' $\beta$ ' represents the standardized coefficients and '*p*' represents significance, with  $p < 0.05$  considered significant (in bold). GFAP, glial fibrillary acidic protein; PET, positron emission tomography;  $^{18}\text{F-SMBT-1}$ , (S)-(2-methylpyrid-5-yl)-6-[(3- $^{18}\text{F}$ -fluoro-2-hydroxy)propoxy] quinoline.

Additionally, age and A $\beta$  PET CL adjusted plasma GFAP ( $\rho = -0.301$ ,  $p = 0.011$ ), and regional  $^{18}\text{F-SMBT-1}$  SUVR (SG:  $\rho = -0.284$ ,  $p = 0.017$ ; PC:  $\rho = -0.301$ ,  $p = 0.011$ ; LT:  $\rho = -0.299$ ,  $p = 0.011$ ; LO:  $\rho = -0.293$ ,  $p = 0.013$ ; and SP:  $\rho = -0.274$ ,  $p = 0.021$ ) were inversely associated with MMSE.

## DISCUSSION

We showed that plasma GFAP was associated with  $^{18}\text{F-SMBT-1}$  PET retention in brain regions that have early A $\beta$  accumulation in the disease trajectory [21], and this relationship appeared to be dependent on levels of insoluble A $\beta$  measured by PET. In line with this observation, significant differences have been reported in both plasma GFAP [7, 12, 14] and  $^{18}\text{F-SMBT-1}$  regional binding [20] between A $\beta$ - and A $\beta$ + participants.  $^{18}\text{F-SMBT-1}$  regional binding is significantly higher in the parietal, temporooccipital, gyrus angularis, posterior cingulate, and supramarginal gyrus regions in AD, and the same regions in addition to orbitofrontal and lateral temporal, also have significantly higher  $^{18}\text{F-SMBT-1}$  binding in HC A $\beta$ + when compared to HC A $\beta$ - [20]. Further, in line with previous studies [6–8, 12, 14], our observations corroborate significant associations between insoluble A $\beta$ , plasma GFAP, and  $^{18}\text{F-SMBT-1}$  PET. We also showed that cognitive performance inversely associated with both plasma GFAP and  $^{18}\text{F-SMBT-1}$  SUVR, independent of age and insoluble A $\beta$ , with similar strength ( $\rho = -0.301$  to  $-0.274$ ).

It has been postulated that the early activation of astrocytes is in response to soluble A $\beta$  oligomers [25, 26] or aggregated A $\beta$  [23, 24]. We found no correlation between  $^{18}\text{F-SMBT-1}$  and plasma levels of soluble A $\beta$  (data not shown); however, this could be attributed to the utilization of plasma monomeric A $\beta$  levels in the current study rather than A $\beta$  oligomer levels. In contrast to the increase in A $\beta$  oligomer levels in plasma of AD patients [44, 45], monomeric A $\beta$  levels decrease in blood plasma [1, 2, 12, 46, 47]. Further, while plasma A $\beta$  likely reflects the soluble biochemical pool of A $\beta$  oligomers in the brain [48], less than 50% of plasma A $\beta$  is derived from A $\beta$  in the brain [49], and this might explain the lack of association with  $^{18}\text{F-SMBT-1}$ . In addition, while MAO-B is overexpressed in GFAP+ astrocytes, especially those surrounding the insoluble A $\beta$  deposits in plaques and cerebral vasculature, not all GFAP+ astrocytes overexpress MAO-B. GFAP expressing astrocytes that

overexpress MAO-B around insoluble A $\beta$  plaques are only a fraction of all GFAP expressing astrocytes, and that might contribute to the observed correlations between plasma GFAP and both insoluble A $\beta$  ( $p < 0.001$ , adjusting for age, sex, and *APOE*  $\epsilon 4$  status) and soluble A $\beta$  ( $p < 0.05$ , adjusting for age, sex, and *APOE*  $\epsilon 4$  status), while  $^{18}\text{F}$ -SMBT-1, assessing MAO-B, was only significantly associated with insoluble A $\beta$  ( $p < 0.005$ , adjusting for age, sex, and *APOE*  $\epsilon 4$  status) but not with soluble A $\beta$  ( $p > 0.05$ , adjusting for age, sex, and *APOE*  $\epsilon 4$  status).

$^{18}\text{F}$ -SMBT-1 is a selective MAO-B tracer with low non-specific binding [19]. In line with the observations in the current study, MAO-B overexpression in reactive astrocytes is significantly associated with GFAP immunoreactivity in AD brain tissue [50, 51]. MAO-B is also upregulated in reactive astrocytes in epilepsy, Parkinson's disease (PD), amyotrophic lateral sclerosis (ALS), and progressive supranuclear palsy (PSP) brain tissue [52–55]. While studies on circulating GFAP levels in other neurological disorders are limited, the current literature indicates significantly higher blood GFAP in epilepsy and PD, but not in ALS and PSP, compared to healthy controls [56–59]. Further, studies report significantly higher GFAP in AD (A $\beta$ +) compared with frontotemporal dementia, dementia with Lewy bodies and healthy controls, and Oeckl and colleagues have suggested that a different type of astrocyte reactivity may be occurring in AD compared to other neurodegenerative diseases [60, 61]. Further studies are required to investigate this aspect in more detail. Whether increased  $^{18}\text{F}$ -SMBT-1 PET signal, reflecting overexpression of MAO-B in reactive astrocytes, can distinguish or be affected by these differences, warrants further investigation.

While MAO-B is mainly overexpressed in the outer membrane of mitochondria in reactive astrocytes [55], it is also expressed in serotonergic neurons [62, 63]. Therefore, it could be posited that the low to moderate strength associations observed between plasma GFAP and  $^{18}\text{F}$ -SMBT-1 PET signal ( $\beta$  with age, sex, and *APOE*  $\epsilon 4$  status adjusted: 0.23 – 0.33) could be partly attributed to cell selectivity. Alternative sensitive and specific neuroimaging and biofluid targets for detecting astrocyte reactivity, particularly those that are also able to distinguish between protective and toxic astrocytic states will aid in understanding the underlying pathophysiology and may provide important information for clinical trials. Additionally, evidence of overexpressed MAO-B as assessed by PET reflecting reactive astrocytes has

been reported in asymptomatic autosomal dominant AD and in prodromal sporadic AD, however a lower PET signal has been reported reaching the dementia phenotype [17, 64]. Additionally, MAO-B PET signal has also been reported to be significantly higher at 6 months than at 8–15 months or 18–24 months in APPswe mice [65]. In contrast, plasma GFAP has been observed to increase along the AD continuum [12, 13]. Together, these observations suggest that the low to moderate strength associations observed between plasma GFAP and  $^{18}\text{F}$ -SMBT-1 PET signal within the current study could also be attributed to the study participant status within the AD continuum.

This study has limitations. The modest sample size of the study might preclude a better evaluation of plasma GFAP relationship with  $^{18}\text{F}$ -SMBT-1 PET across the AD continuum (HC A $\beta$ +, MCI A $\beta$ +, and AD A $\beta$ +) . Analyses within the subset of participants in the AD continuum within the current study had no significant association between plasma GFAP and SMBT-1 PET (Supplementary Table 3A), but within the cognitively impaired individual subset (MCI and AD participants), a significant or trend towards significant association was found between plasma GFAP and SMBT-1 PET (Supplementary Table 3B), which could be attributed to the latter subset comprising both A $\beta$ + and A $\beta$ - individuals. This suggests that the association between plasma GFAP and SMBT-1 is dependent on brain A $\beta$  load or A $\beta$  status. However, the small sample size being a limitation of the study is acknowledged and further confirmatory studies are required. In addition, the assessments were cross-sectional in nature, and it will be necessary to compare and contrast these findings with longitudinal trajectories to determine changes over time and/or disease progression. Further, fifty-nine participants of the 71 participants in this study, underwent  $^{18}\text{F}$ -SMBT-1 imaging within 12 months of blood collection, while 12 participants had  $^{18}\text{F}$ -SMBT-1 imaging over 12 months from blood collection (analyses for this subset has been provided in Supplementary Tables 2 and 4). Further, participants from the study cohort were volunteers and not randomly selected from the community, and due to the strict inclusion/exclusion criteria the results may not be applicable to the general population. Additionally, this study did not include other neurodegenerative conditions, although it is acknowledged that neuroinflammation is not disease-specific and astrocyte activity markers are associated with various conditions including traumatic brain injury, major depressive disorder, and several neurodegen-

erative diseases [66–70]. Further studies also need to investigate whether there is an association between CSF GFAP and  $^{18}\text{F}$ -SMBT-1 PET, and if it is also dependent on A $\beta$  load. Unlike plasma GFAP, CSF GFAP is not elevated in cognitively unimpaired A $\beta$ + individuals when compared to cognitively unimpaired A $\beta$ - individuals [13]. Such investigations will also tell us whether  $^{18}\text{F}$ -SMBT-1 is more specific to AD, given that MAO-B is overexpressed in GFAP+ astrocytes surrounding the insoluble A $\beta$  plaques.

To conclude, an association between the astrocyte markers, plasma GFAP and regional  $^{18}\text{F}$ -SMBT-1 PET, was observed. However, this relationship appears to be dependent on brain A $\beta$  load. Follow-up studies assessing how this relationship evolves over time are ongoing.

## ACKNOWLEDGMENTS

We thank all the participants and their families who took part in this study as well as the clinicians who referred participants. The AIBL study (<http://www.AIBL.csiro.au>) is a collaboration between CSIRO, Edith Cowan University (ECU), National Ageing Research Institute (NARI), The Florey Institute of Neuroscience and Mental Health (FINMH) and Austin Health. The study also received support from Hollywood Private Hospital, Sir Charles Gairdner Hospital, CSIRO, the Science and Industry Endowment Fund, and the Australian Alzheimer's Research Foundation. The authors acknowledge the financial support of the Cooperative Research Centre (CRC) for Mental Health, an Australian Government Initiative. Pfizer International has provided financial support to assist with analysis of blood samples and to further the AIBL research program. We are grateful to the Lions Alzheimer's Foundation and Lions Club International for their generous donations that allowed the purchase of the Simoa-HD-X instrument used in this study.

## FUNDING

Financial support was received from the Cooperative Research Centre (CRC) for Mental Health, an Australian Government Initiative. Pfizer International has provided financial support to assist with analysis of blood samples and to further the AIBL research program. The Lions Alzheimer's Foundation and Lions Club International provided donations that allowed the purchase of the Simoa-HD-X instru-

ment used in this study. PC is funded by Macquarie University and RNM is jointly funded by Macquarie University and Edith Cowan University.

## CONFLICT OF INTEREST

SRS has received grant support from the National Health and Medical Research Council, Alzheimer's Association (USA) Research Grant, Alzheimer's Drug Discovery Foundation and the BrightFocus Foundation and honorarium for lectures from the Mature Adults Learning Association Inc. SCB is a current employee and minor stockholder of Eli Lilly and Company. KT and RNM are Directors of SMarT Minds Western Australia. CCR has received research grants from NHMRC, Enigma Australia, Biogen, Eisai and Abbvie. He is on the scientific advisory board for Cerveau Technologies and consulted for Prothena, Eisai, Roche and Biogen Australia. VLV has received research grants from NHMRC (GNT2001320), the Aging Mind Foundation (DAF2255207) and NIH 2P01AG025204-16) and is and has been a consultant or paid speaker at sponsored conference sessions for Eli Lilly, Life Molecular Imaging, ACE Barcelona, and IXICO. Colin L. Masters and Ralph N. Martins are Editorial Board Members of this journal but were not involved in the peer-review process nor had access to any information regarding its peer-review. The other authors did not report any conflict of interest.

## DATA AVAILABILITY

The data supporting the findings of this study are available on request. The data are not publicly available due to privacy and ethical restrictions.

## SUPPLEMENTARY MATERIAL

The supplementary material is available in the electronic version of this article: <https://dx.doi.org/10.3233/JAD-220908>.

## REFERENCES

- [1] Nakamura A, Kaneko N, Villemagne VL, Kato T, Doecke J, Dore V, Fowler C, Li QX, Martins R, Rowe C, Tomita T, Matsuzaki K, Ishii K, Ishii K, Arahata Y, Iwamoto S, Ito K, Tanaka K, Masters CL, Yanagisawa K (2018) High performance plasma amyloid-beta biomarkers for Alzheimer's disease. *Nature* **554**, 249-254.
- [2] Ovod V, Ramsey KN, Mawuenyega KG, Bollinger JG, Hicks T, Schneider T, Sullivan M, Paumier K, Holtzman

- DM, Morris JC, Benzinger T, Fagan AM, Patterson BW, Bateman RJ (2017) Amyloid beta concentrations and stable isotope labeling kinetics of human plasma specific to central nervous system amyloidosis. *Alzheimers Dement* **13**, 841-849.
- [3] Karikari TK, Pascoal TA, Ashton NJ, Janelidze S, Benedet AL, Rodriguez JL, Chamoun M, Savard M, Kang MS, Theriault J, Scholl M, Massarweh G, Soucy JP, Hoglund K, Brinkmalm G, Mattsson N, Palmqvist S, Gauthier S, Stomrud E, Zetterberg H, Hansson O, Rosa-Neto P, Blennow K (2020) Blood phosphorylated tau 181 as a biomarker for Alzheimer's disease: A diagnostic performance and prediction modelling study using data from four prospective cohorts. *Lancet Neurol* **19**, 422-433.
- [4] Palmqvist S, Janelidze S, Quiroz YT, Zetterberg H, Lopera F, Stomrud E, Su Y, Chen Y, Serrano GE, Leuzy A, Mattsson-Carlsson N, Strandberg O, Smith R, Villegas A, Sepulveda-Falla D, Chai X, Proctor NK, Beach TG, Blennow K, Dage JL, Reiman EM, Hansson O (2020) Discriminative accuracy of plasma phospho-tau217 for Alzheimer disease vs other neurodegenerative disorders. *JAMA* **324**, 772-781.
- [5] Ashton NJ, Pascoal TA, Karikari TK, Benedet AL, Lantero-Rodriguez J, Brinkmalm G, Snellman A, Scholl M, Troakes C, Hye A, Gauthier S, Vanmechelen E, Zetterberg H, Rosa-Neto P, Blennow K (2021) Plasma p-tau231: A new biomarker for incipient Alzheimer's disease pathology. *Acta Neuropathol* **141**, 709-724.
- [6] Verberk IMW, Thijssen E, Koelewijn J, Mauroo K, Vanbrabant J, de Wilde A, Zwan MD, Verfaillie SCJ, Ossenkoppele R, Barkhof F, van Berckel BNM, Scheltens P, van der Flier WM, Stoops E, Vanderstichele HM, Teunissen CE (2020) Combination of plasma amyloid beta(1-42/1-40) and glial fibrillary acidic protein strongly associates with cerebral amyloid pathology. *Alzheimers Res Ther* **12**, 118.
- [7] Chatterjee P, Pedrini S, Stoops E, Goozee K, Villemagne VL, Asih PR, Verberk IMW, Dave P, Taddei K, Sohrabi HR, Zetterberg H, Blennow K, Teunissen CE, Vanderstichele HM, Martins RN (2021) Plasma glial fibrillary acidic protein is elevated in cognitively normal older adults at risk of Alzheimer's disease. *Transl Psychiatry* **11**, 27.
- [8] Pereira JB, Janelidze S, Smith R, Mattsson-Carlsson N, Palmqvist S, Teunissen CE, Zetterberg H, Stomrud E, Ashton NJ, Blennow K, Hansson O (2021) Plasma GFAP is an early marker of amyloid-beta but not tau pathology in Alzheimer's disease. *Brain* **144**, 3505-3516.
- [9] Escartin C, Galea E, Lakatos A, O'Callaghan JP, Petzold GC, Serrano-Pozo A, Steinhilber C, Volterra A, Carmignoto G, Agarwal A, Allen NJ, Araque A, Barbeito L, Barzilai A, Bergles DE, Bonvento G, Butt AM, Chen WT, Cohen-Salmon M, Cunningham C, Deneen B, De Strooper B, Diaz-Castro B, Farina C, Freeman M, Gallo V, Goldman JE, Goldman SA, Gotz M, Gutierrez A, Haydon PG, Heiland DH, Hol EM, Holt MG, Iino M, Kastanenka KV, Kettenmann H, Khakh BS, Koizumi S, Lee CJ, Lidde-low SA, MacVicar BA, Magistretti P, Messing A, Mishra A, Molofsky AV, Murai KK, Norris CM, Okada S, Oliet SHR, Oliveira JF, Panatier A, Parpura V, Pekna M, Pekny M, Pellerin L, Perea G, Perez-Nievas BG, Pfrieger FW, Poskanzer KE, Quintana FJ, Ransohoff RM, Riquelme-Perez M, Robel S, Rose CR, Rothstein JD, Rouach N, Rowitch DH, Semyanov A, Sirko S, Sontheimer H, Swanson RA, Vitorica J, Wanner IB, Wood LB, Wu J, Zheng B, Zimmer ER, Zorec R, Sofroniew MV, Verkhratsky A (2021) Reactive astrocyte nomenclature, definitions, and future directions. *Nat Neurosci* **24**, 312-325.
- [10] Muramori F, Kobayashi K, Nakamura I (1998) A quantitative study of neurofibrillary tangles, senile plaques and astrocytes in the hippocampal subdivisions and entorhinal cortex in Alzheimer's disease, normal controls and non-Alzheimer neuropsychiatric diseases. *Psychiatry Clin Neurosci* **52**, 593-599.
- [11] Kamphuis W, Middeldorp J, Kooijman L, Sluijs JA, Kooi EJ, Moeton M, Freriks M, Mizee MR, Hol EM (2014) Glial fibrillary acidic protein isoform expression in plaque related astrogliosis in Alzheimer's disease. *Neurobiol Aging* **35**, 492-510.
- [12] Chatterjee P, Pedrini S, Doecke JD, Thota R, Villemagne VL, Doré V, Singh AK, Wang P, Rainey-Smith S, Fowler C, Taddei K, Sohrabi HR, Molloy MP, Ames D, Maruff P, Rowe CC, Masters CL, Martins RN (2022) Plasma A $\beta$ 42/40 ratio, p-tau181, GFAP and NFL across the Alzheimer's disease continuum: A cross-sectional and longitudinal study in the AIBL cohort. *Alzheimers Dement*. doi: 10.1002/alz.12724.
- [13] Benedet AL, Milà-Alomà M, Vrillon A, Ashton NJ, Pascoal TA, Lussier F, Karikari TK, Hourregue C, Cognat E, Dumurgier J, Stevenson J, Rahmouni N, Pallen V, Poltronetti NM, Salvadó G, Shekari M, Operto G, Gispert JD, Minguillon C, Fauria K, Kollmorgen G, Suridjan I, Zimmer ER, Zetterberg H, Molinuevo JL, Paquet C, Rosa-Neto P, Blennow K, Suárez-Calvet M; Translational Biomarkers in Aging and Dementia (TRIAD) study, Alzheimer's and Families (ALFA) study, and BioCogBank Paris Lariboisière cohort (2021) Differences between plasma and cerebrospinal fluid glial fibrillary acidic protein levels across the Alzheimer disease continuum. *JAMA Neurol* **78**, 1471-1483.
- [14] Chatterjee P, Pedrini S, Ashton NJ, Tegg M, Goozee K, Singh AK, Karikari TK, Simren J, Vanmechelen E, Armstrong NJ, Hone E, Asih PR, Taddei K, Dore V, Villemagne VL, Sohrabi HR, Zetterberg H, Masters CL, Blennow K, Martins RN (2022) Diagnostic and prognostic plasma biomarkers for preclinical Alzheimer's disease. *Alzheimers Dement* **18**, 1141-1154.
- [15] Cicognola C, Janelidze S, Hertze J, Zetterberg H, Blennow K, Mattsson-Carlsson N, Hansson O (2021) Plasma glial fibrillary acidic protein detects Alzheimer pathology and predicts future conversion to Alzheimer dementia in patients with mild cognitive impairment. *Alzheimers Res Ther* **13**, 68.
- [16] Carter SF, Scholl M, Almkvist O, Wall A, Engler H, Langstrom B, Nordberg A (2012) Evidence for astrocytosis in prodromal Alzheimer disease provided by 11C-deuterium-L-deprenyl: A multitracers PET paradigm combining 11C-Pittsburgh compound B and 18F-FDG. *J Nucl Med* **53**, 37-46.
- [17] Rodriguez-Vieitez E, Saint-Aubert L, Carter SF, Almkvist O, Farid K, Scholl M, Chiotis K, Thordardottir S, Graff C, Wall A, Langstrom B, Nordberg A (2016) Diverging longitudinal changes in astrocytosis and amyloid PET in autosomal dominant Alzheimer's disease. *Brain* **139**, 922-936.
- [18] Harada R, Hayakawa Y, Ezura M, Lerdsiriruk P, Du Y, Ishikawa Y, Iwata R, Shidahara M, Ishiki A, Kikuchi A, Arai H, Kudo Y, Yanai K, Furumoto S, Okamura N (2021) (18)F-SMBT-1: A selective and reversible PET tracer for monoamine oxidase-B imaging. *J Nucl Med* **62**, 253-258.
- [19] Villemagne VL, Harada R, Dore V, Furumoto S, Mulligan R, Kudo Y, Burnham S, Krishnadas N, Bozinovsk

- S, Huang K, Lopresti BJ, Yanai K, Rowe CC, Okamura N (2022) First-in-humans evaluation of (18)F-SMT-1, a novel (18)F-labeled monoamine oxidase-B PET tracer for imaging reactive astrogliosis. *J Nucl Med* **63**, 1551-1559.
- [20] Villemagne VL, Harada R, Dore V, Furumoto S, Mulligan R, Kudo Y, Burnham S, Krishnadas N, Bourgeat P, Xia Y, Laws S, Bozinovski S, Huang K, Ikonovic MD, Frapp J, Yanai K, Okamura N, Rowe CC (2022) Assessing reactive astrogliosis with (18)F-SMT-1 across the Alzheimer disease spectrum. *J Nucl Med* **63**, 1560-1569.
- [21] Villain N, Chetelat G, Grassiot B, Bourgeat P, Jones G, Ellis KA, Ames D, Martins RN, Eustache F, Salvado O, Masters CL, Rowe CC, Villemagne VL, Group AR (2012) Regional dynamics of amyloid-beta deposition in healthy elderly, mild cognitive impairment and Alzheimer's disease: A voxelwise PiB-PET longitudinal study. *Brain* **135**, 2126-2139.
- [22] Guo T, Landau SM, Jagust WJ, Alzheimer's Disease Neuroimaging Initiative (2020) Detecting earlier stages of amyloid deposition using PET in cognitively normal elderly adults. *Neurology* **94**, e1512-e1524.
- [23] Hu J, Akama KT, Krafft GA, Chromy BA, Van Eldik LJ (1998) Amyloid-beta peptide activates cultured astrocytes: Morphological alterations, cytokine induction and nitric oxide release. *Brain Res* **785**, 195-206.
- [24] Meda L, Baron P, Scarlato G (2001) Glial activation in Alzheimer's disease: The role of Abeta and its associated proteins. *Neurobiol Aging* **22**, 885-893.
- [25] Wyssbach A, Quintela T, Llavero F, Zugaza JL, Matute C, Alberdi E (2016) Amyloid beta-induced astrogliosis is mediated by beta1-integrin via NADPH oxidase 2 in Alzheimer's disease. *Aging Cell* **15**, 1140-1152.
- [26] Forloni G, Balducci C (2018) Alzheimer's disease, oligomers, and inflammation. *J Alzheimers Dis* **62**, 1261-1276.
- [27] Chatterjee P, Vermunt L, Gordon BA, Pedrini S, Boonkamp L, Armstrong NJ, Xiong C, Singh AK, Li Y, Sohrabi HR, Taddei K, Molloy M, Benzinger TLS, Morris JC, Karch C, Berman S, Chhatwal J, Cruchaga C, Graff-Radford NR, Day GS, Farlow M, Fox N, Goate A, Hassenstab J, Lee JH, Levin J, McDade E, Mori H, Perrin R, Sanchez-Valle R, Schofield PR, Levey A, Jucker M, Masters CL, Fagan AM, Bateman RJ, Martins RN, Teunissen C, and the Dominantly Inherited Alzheimer Network (2022) Plasma glial fibrillary acidic protein in autosomal dominant Alzheimer's disease: Associations with Abeta-PET, neurodegeneration, and cognition. *Alzheimers Dement*. doi: 10.1002/alz.12879.
- [28] Ellis KA, Bush AI, Darby D, De Fazio D, Foster J, Hudson P, Lautenschlager NT, Lenzo N, Martins RN, Maruff P, Masters C, Milner A, Pike K, Rowe C, Savage G, Szeoke C, Taddei K, Villemagne V, Woodward M, Ames D, AIBL Research Group (2009) The Australian Imaging, Biomarkers and Lifestyle (AIBL) study of aging: Methodology and baseline characteristics of 1112 individuals recruited for a longitudinal study of Alzheimer's disease. *Int Psychogeriatr* **21**, 672-687.
- [29] Fowler C, Rainey-Smith SR, Bird S, Bomke J, Bourgeat P, Brown BM, Burnham SC, Bush AI, Chadunow C, Collins S, Doecke J, Doré V, Ellis KA, Evered L, Fazlollahi A, Frapp J, Gardener SL, Gibson S, Grenfell R, Harrison E, Head R, Jin L, Kamer A, Lamb F, Lautenschlager NT, Laws SM, Li QX, Lim L, Lim YY, Louey A, Macaulay SL, Mackintosh L, Martins RN, Maruff P, Masters CL, McBride S, Milicich L, Peretti M, Pertile K, Porter T, Radler M, Rembach A, Robertson J, Rodrigues M, Rowe CC, Rumble R, Salvado O, Savage G, Silbert B, Soh M, Sohrabi HR, Taddei K, Taddei T, Thai C, Trounson B, Tyrrell R, Vacher M, Varghese S, Villemagne VL, Weinborn M, Woodward M, Xia Y, Ames D; the AIBL investigators (2021) Fifteen years of the Australian Imaging, Biomarkers and Lifestyle (AIBL) Study: Progress and observations from 2,359 older adults spanning the spectrum from cognitive normality to Alzheimer's disease. *J Alzheimers Dis Rep* **5**, 443-468.
- [30] McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM (1984) Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* **34**, 939-944.
- [31] Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E (1999) Mild cognitive impairment: Clinical characterization and outcome. *Arch Neurol* **56**, 303-308.
- [32] Winblad B, Palmer K, Kivipelto M, Jelic V, Fratiglioni L, Wahlund LO, Nordberg A, Backman L, Albert M, Almkvist O, Arai H, Basun H, Blennow K, de Leon M, DeCarli C, Erkinjuntti T, Giacobini E, Graff C, Hardy J, Jack C, Jorm A, Ritchie K, van Duijn C, Visser P, Petersen RC (2004) Mild cognitive impairment—beyond controversies, towards a consensus: Report of the International Working Group on Mild Cognitive Impairment. *J Intern Med* **256**, 240-246.
- [33] Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* **12**, 189-198.
- [34] Morris JC (1993) The Clinical Dementia Rating (CDR): Current version and scoring rules. *Neurology* **43**, 2412-2414.
- [35] Battle MR, Pillay LC, Lowe VJ, Knopman D, Kemp B, Rowe CC, Dore V, Villemagne VL, Buckley CJ (2018) Centiloid scaling for quantification of brain amyloid with [(18)F]flutemetamol using multiple processing methods. *EJNMMI Res* **8**, 107.
- [36] Rowe CC, Jones G, Dore V, Pejaska S, Margison L, Mulligan RS, Chan JG, Young K, Villemagne VL (2016) Standardized expression of 18F-NAV4694 and 11C-PiB beta-amyloid PET results with the Centiloid scale. *J Nucl Med* **57**, 1233-1237.
- [37] Navitsky M, Joshi AD, Kennedy I, Klunk WE, Rowe CC, Wong DF, Pontecorvo MJ, Mintun MA, Devous MD, Sr. (2018) Standardization of amyloid quantitation with florbetapir standardized uptake value ratios to the Centiloid scale. *Alzheimers Dement* **14**, 1565-1571.
- [38] Klunk WE, Koeppe RA, Price JC, Benzinger TL, Devous MD, Sr., Jagust WJ, Johnson KA, Mathis CA, Minhas D, Pontecorvo MJ, Rowe CC, Skovronsky DM, Mintun MA (2015) The Centiloid Project: Standardizing quantitative amyloid plaque estimation by PET. *Alzheimers Dement* **11**, 1-15 e11-14.
- [39] Bourgeat P, Dore V, Frapp J, Ames D, Masters CL, Salvado O, Villemagne VL, Rowe CC, AIBL research group (2018) Implementing the centiloid transformation for (11)C-PiB and beta-amyloid (18)F-PET tracers using CapAIBL. *Neuroimage* **183**, 387-393.
- [40] La Joie R, Ayakta N, Seeley WW, Borys E, Boxer AL, DeCarli C, Dore V, Grinberg LT, Huang E, Hwang JH, Ikonovic MD, Jack C Jr, Jagust WJ, Jin LW, Klunk WE, Kofler J, Lesman-Segev OH, Lockhart SN, Lowe VJ, Masters CL, Mathis CA, McLean CL, Miller BL, Mungas D, O'Neil JP, Olichney JM, Parisi JE, Petersen RC, Rosen HJ, Rowe CC, Spina S, Vemuri P, Villemagne VL, Murray ME, Rabinovici GD (2019) Multisite study of the relationships between antemortem [(11)C]PIB-PET Centiloid values and

- postmortem measures of Alzheimer's disease neuropathology. *Alzheimers Dement* **15**, 205-216.
- [41] Jack CR, Jr., Wiste HJ, Weigand SD, Therneau TM, Lowe VJ, Knopman DS, Gunter JL, Senjem ML, Jones DT, Kantarci K, Machulda MM, Mielke MM, Roberts RO, Vemuri P, Reyes DA, Petersen RC (2017) Defining imaging biomarker cut points for brain aging and Alzheimer's disease. *Alzheimers Dement* **13**, 205-216.
- [42] Zhou L, Salvado O, Dore V, Bourgeat P, Raniga P, Macaulay SL, Ames D, Masters CL, Ellis KA, Villemagne VL, Rowe CC, Frupp J, AIBL Research Group (2014) MR-less surface-based amyloid assessment based on 11C PiB PET. *PLoS One* **9**, e84777.
- [43] Bourgeat P, Villemagne VL, Dore V, Brown B, Macaulay SL, Martins R, Masters CL, Ames D, Ellis K, Rowe CC, Salvado O, Frupp J, Group AR (2015) Comparison of MR-less PiB SUVR quantification methods. *Neurobiol Aging* **36 Suppl 1**, S159-166.
- [44] Wang MJ, Yi S, Han JY, Park SY, Jang JW, Chun IK, Kim SE, Lee BS, Kim GJ, Yu JS, Lim K, Kang SM, Park YH, Youn YC, An SSA, Kim S (2017) Oligomeric forms of amyloid-beta protein in plasma as a potential blood-based biomarker for Alzheimer's disease. *Alzheimers Res Ther* **9**, 98.
- [45] Babapour Mofrad R, Scheltens P, Kim S, Kang S, Youn YC, An SSA, Tomassen J, van Berckel BNM, Visser PJ, van der Flier WM, Teunissen CE (2021) Plasma amyloid-beta oligomerization assay as a pre-screening test for amyloid status. *Alzheimers Res Ther* **13**, 133.
- [46] Janelidze S, Stomrud E, Palmqvist S, Zetterberg H, van Westen D, Jeromin A, Song L, Hanlon D, Tan Hehir CA, Baker D, Blennow K, Hansson O (2016) Plasma beta-amyloid in Alzheimer's disease and vascular disease. *Sci Rep* **6**, 26801.
- [47] Verberk IMW, Slot RE, Verfaillie SCJ, Heijst H, Prins ND, van Berckel BNM, Scheltens P, Teunissen CE, van der Flier WM (2018) Plasma amyloid as prescreener for the earliest Alzheimer pathological changes. *Ann Neurol* **84**, 648-658.
- [48] Roberts BR, Lind M, Wagen AZ, Rembach A, Frugier T, Li QX, Ryan TM, McLean CA, Doecke JD, Rowe CC, Villemagne VL, Masters CL (2017) Biochemically-defined pools of amyloid-beta in sporadic Alzheimer's disease: Correlation with amyloid PET. *Brain* **140**, 1486-1498.
- [49] Roberts KF, Elbert DL, Kasten TP, Patterson BW, Sigurdson WC, Connors RE, Ovod V, Munsell LY, Mawuenyega KG, Miller-Thomas MM, Moran CJ, Cross DT, 3rd, Derdeyn CP, Bateman RJ (2014) Amyloid-beta efflux from the central nervous system into the plasma. *Ann Neurol* **76**, 837-844.
- [50] Saura J, Luque JM, Cesura AM, Da Prada M, Chan-Palay V, Huber G, Loffler J, Richards JG (1994) Increased monoamine oxidase B activity in plaque-associated astrocytes of Alzheimer brains revealed by quantitative enzyme radioautography. *Neuroscience* **62**, 15-30.
- [51] Jo S, Yarishkin O, Hwang YJ, Chun YE, Park M, Woo DH, Bae JY, Kim T, Lee J, Chun H, Park HJ, Lee DY, Hong J, Kim HY, Oh SJ, Park SJ, Lee H, Yoon BE, Kim Y, Jeong Y, Shim I, Bae YC, Cho J, Kowall NW, Ryu H, Hwang E, Kim D, Lee CJ (2014) GABA from reactive astrocytes impairs memory in mouse models of Alzheimer's disease. *Nat Med* **20**, 886-896.
- [52] Kumlien E, Hilton-Brown P, Spannare B, Gillberg PG (1992) In vitro quantitative autoradiography of [3H]-L-deprenyl and [3H]-PK 11195 binding sites in human epileptic hippocampus. *Epilepsia* **33**, 610-617.
- [53] Tong J, Rathitharan G, Meyer JH, Furukawa Y, Ang LC, Boileau I, Guttman M, Hornykiewicz O, Kish SJ (2017) Brain monoamine oxidase B and A in human parkinsonian dopamine deficiency disorders. *Brain* **140**, 2460-2474.
- [54] Aquilonius SM, Jossan SS, Ekblom JG, Askmark H, Gillberg PG (1992) Increased binding of 3H-L-deprenyl in spinal cords from patients with amyotrophic lateral sclerosis as demonstrated by autoradiography. *J Neural Transm Gen Sect* **89**, 111-122.
- [55] Ekblom J, Jossan SS, Bergstrom M, Orelund L, Walum E, Aquilonius SM (1993) Monoamine oxidase-B in astrocytes. *Glia* **8**, 122-132.
- [56] Simani L, Elmi M, Asadollahi M (2018) Serum GFAP level: A novel adjunctive diagnostic test in differentiate epileptic seizures from psychogenic attacks. *Seizure* **61**, 41-44.
- [57] Su W, Chen HB, Li SH, Wu DY (2012) Correlational study of the serum levels of the glial fibrillary acidic protein and neurofilament proteins in Parkinson's disease patients. *Clin Neurol Neurosurg* **114**, 372-375.
- [58] Oeckl P, Weydt P, Steinacker P, Anderl-Straub S, Nordin F, Volk AE, Diehl-Schmid J, Andersen PM, Kornhuber J, Danek A, Fassbender K, Fliessbach K, German Consortium for Frontotemporal Lobar D, Jahn H, Lauer M, Muller K, Knehr A, Prudlo J, Schneider A, Thal DR, Yilmazer-Hanke D, Weishaupt JH, Ludolph AC, Otto M (2019) Different neuroinflammatory profile in amyotrophic lateral sclerosis and frontotemporal dementia is linked to the clinical phase. *J Neurol Neurosurg Psychiatry* **90**, 4-10.
- [59] Benussi A, Ashton NJ, Karikari TK, Gazzina S, Premi E, Benussi L, Ghidoni R, Rodriguez JL, Emersic A, Binetti G, Fostinelli S, Giunta M, Gasparotti R, Zetterberg H, Blennow K, Borroni B (2020) Serum glial fibrillary acidic protein (GFAP) is a marker of disease severity in frontotemporal lobar degeneration. *J Alzheimers Dis* **77**, 1129-1141.
- [60] Thijssen EH, Verberk IMW, Kindermans J, Abramian A, Vanbrabant J, Ball AJ, Pijnenburg Y, Lemstra AW, van der Flier WM, Stoops E, Hirtz C, Teunissen CE (2022) Differential diagnostic performance of a panel of plasma biomarkers for different types of dementia. *Alzheimers Dement (Amst)* **14**, e12285.
- [61] Oeckl P, Anderl-Straub S, Von Arnim CAF, Baldeiras I, Diehl-Schmid J, Grimmer T, Halbgebauer S, Kort AM, Lima M, Marques TM, Ortner M, Santana I, Steinacker P, Verbeeck MM, Volk AE, Ludolph AC, Otto M (2022) Serum GFAP differentiates Alzheimer's disease from frontotemporal dementia and predicts MCI-to-dementia conversion. *J Neurol Neurosurg Psychiatry*. doi: 10.1136/jnnp-2021-328547.
- [62] Levitt P, Pintar JE, Breakefield XO (1982) Immunocytochemical demonstration of monoamine oxidase B in brain astrocytes and serotonergic neurons. *Proc Natl Acad Sci U S A* **79**, 6385-6389.
- [63] Harada R, Furumoto S, Kudo Y, Yanai K, Villemagne VL, Okamura N (2022) Imaging of reactive astrogliosis by positron emission tomography. *Front Neurosci* **16**, 807435.
- [64] Vilaplana E, Rodriguez-Veitez E, Ferreira D, Montal V, Almkvist O, Wall A, Lleo A, Westman E, Graff C, Fortea J, Nordberg A (2020) Cortical microstructural correlates of astrogliosis in autosomal-dominant Alzheimer disease. *Neurology* **94**, e2026-e2036.
- [65] Rodriguez-Veitez E, Ni R, Gulyas B, Toth M, Haggkvist J, Halldin C, Voytenko L, Marutle A, Nordberg A (2015) Astrogliosis precedes amyloid plaque deposition in Alzheimer APPsw transgenic mouse brain: A correlative

- positron emission tomography and in vitro imaging study. *Eur J Nucl Med Mol Imaging* **42**, 1119-1132.
- [66] Yue JK, Yuh EL, Korley FK, Winkler EA, Sun X, Puffer RC, Deng H, Choy W, Chandra A, Taylor SR, Ferguson AR, Huie JR, Rabinowitz M, Puccio AM, Mukherjee P, Vassar MJ, Wang KKW, Diaz-Arrastia R, Okonkwo DO, Jain S, Manley GT, Investigators T-T (2019) Association between plasma GFAP concentrations and MRI abnormalities in patients with CT-negative traumatic brain injury in the TRACK-TBI cohort: A prospective multicentre study. *Lancet Neurol* **18**, 953-961.
- [67] Huebschmann NA, Luoto TM, Karr JE, Berghem K, Blennow K, Zetterberg H, Ashton NJ, Simren J, Posti JP, Gill JM, Iverson GL (2020) Comparing glial fibrillary acidic protein (GFAP) in serum and plasma following mild traumatic brain injury in older adults. *Front Neurol* **11**, 1054.
- [68] Michel M, Fiebich BL, Kuzior H, Meixensberger S, Berger B, Maier S, Nickel K, Runge K, Denzel D, Pankratz B, Schiele MA, Domschke K, van Elst LT, Endres D (2021) Increased GFAP concentrations in the cerebrospinal fluid of patients with unipolar depression. *Transl Psychiatry* **11**, 308.
- [69] Steinacker P, Al Shweiki MR, Oeckl P, Graf H, Ludolph AC, Schonfeldt-Lecuona C, Otto M (2021) Glial fibrillary acidic protein as blood biomarker for differential diagnosis and severity of major depressive disorder. *J Psychiatr Res* **144**, 54-58.
- [70] Baiardi S, Quadalti C, Mammama A, Dellavalle S, Zenesini C, Sambati L, Pantieri R, Polisch B, Romano L, Suffritti M, Bentivenga GM, Randi V, Stanzani-Maserati M, Capellari S, Parchi P (2022) Diagnostic value of plasma p-tau181, NFL, and GFAP in a clinical setting cohort of prevalent neurodegenerative dementias. *Alzheimers Res Ther* **14**, 153.