

# Supplementary Material

## Geographic Barriers Drive Disparities in Specialty Center Access for Older Adults with Huntington's Disease

**Supplementary Table 1.** State level variation in proportion of Medicare beneficiaries with HD proximal to HDSA COE, 2017

State	0-100 miles (%)*	% 101+ miles (%)
Alabama	57.30%	42.70%
Alaska	0.00%	100.00%
Arizona	0.00%	100.00%
Arkansas	14.70%	85.30%
California	78.30%	21.70%
Colorado	84.80%	15.20%
Connecticut	65.60%	34.40%
Delaware	80.00%	20.00%
District of Columbia	100.00%	0.00%
Florida	76.70%	23.30%
Georgia	61.80%	38.20%
Guam	0.00%	100.00%
Hawaii	0.00%	100.00%
Idaho	2.60%	97.40%
Illinois	80.40%	19.60%
Indiana	81.80%	18.20%
Iowa	56.50%	43.50%
Kansas	0.00%	100.00%
Kentucky	65.90%	34.10%
Louisiana	40.40%	59.60%
Maine	5.00%	95.00%
Maryland	93.80%	6.20%
Massachusetts	98.90%	1.10%
Michigan	0.90%	99.10%
Minnesota	69.70%	30.30%
Mississippi	18.30%	81.70%
Missouri	41.20%	58.80%
Montana	0.00%	100.00%
Nebraska	56.50%	43.50%
Nevada	0.00%	100.00%
New Hampshire	93.80%	6.20%
New Jersey	99.60%	0.40%
New Mexico	0.00%	100.00%
New York	92.60%	7.40%
North Carolina	44.80%	55.20%

North Dakota	0.00%	100.00%
Ohio	78.80%	21.20%
Oklahoma	0.00%	100.00%
Oregon	0.00%	100.00%
Pennsylvania	75.20%	24.80%
Puerto Rico	0.00%	100.00%
Rhode Island	100.00%	0.00%
South Carolina	0.00%	100.00%
South Dakota	0.00%	100.00%
Tennessee	60.20%	39.80%
Texas	27.70%	72.30%
Utah	87.00%	13.00%
Vermont	97.40%	2.60%
Virgin Islands	0.00%	100.00%
Virginia	67.60%	32.40%
Washington	67.50%	32.50%
West Virginia	17.60%	82.40%
Wisconsin	54.80%	45.20%
Wyoming	0.00%	100.00%

\*Medicare Data Use Agreement prevents reporting of small cell sizes, therefore only percentages data is provided.