

Supplementary Material

Advancing Parkinson's Disease Research in Canada: The Canadian Open Parkinson Network (C-OPN) Cohort

Supplementary Table 1. Differences in key features by region. Western Canada (British Columbia and Alberta) relative to eastern Canada (Ontario and Quebec).

West Canada*	Odds ratio	Error	z	p	Low CI	High CI
Dyskinesias	1.503	0.240	2.550	0.011	1.099	2.054
Deglutition therapy	3.668	2.232	2.140	0.033	1.113	12.088
Instability at onset	1.759	0.474	2.100	0.036	1.038	2.981
DBS in subthalamus	6.460	4.444	2.710	0.007	1.678	24.876
Pramipexol	0.312	0.074	-4.930	0.000	0.196	0.495
Levodopa/carbidopa/entacapone	0.149	0.061	-4.650	0.000	0.067	0.333
PD in brothers	0.344	0.182	-2.020	0.043	0.122	0.969
Constant	0.965	0.105	-0.330	0.744	0.779	1.195

West Canada†	Odds ratio	Error	z	p	Low CI	High CI
DBS in subthalamus	6.282	3.474	3.320	0.001	2.125	18.571
Age at onset (years, z)	1.226	0.081	3.100	0.002	1.078	1.395
PD in brothers	0.370	0.162	-2.270	0.023	0.157	0.872
Constant	1.177	0.077	2.500	0.012	1.036	1.337

* Fitted with variables having less than 33% missing values.

† Fitted with variables having less than 20% missing values.