

## Erratum

---

### Neuromuscular disorders in children in South-Eastern Norway

Magnhild Rasmussen, Knut Risberg, Arve Vøllo and Ola H. Skjeldal

[*Journal of Pediatric Neurology* 10(2) (2012), 95–100]

When this paper was originally published, Table 2 was displayed incomplete. The correct Table 2 is presented below. The original article can be found here:

<http://iospress.metapress.com/content/t133672503501647/?p=85a35a280af3476ca2a8f442603652ff&pi=2>.

Table 2  
Prevalence figures for some neuromuscular disorders in the investigated population

| Diagnosis                        | Prevalence ( $\times 10^{-5}$ ) |
|----------------------------------|---------------------------------|
| Duchenne muscular dystrophy      | 16.2 <sup>a,b</sup>             |
| Becker muscular dystrophy        | 3.5 <sup>a</sup>                |
| Peripheral neuropathies          | 8.1                             |
| Myotonic dystrophy               | 3.3                             |
| Spinal muscular atrophy          | 2.5                             |
| Limb girdle muscular dystrophies | 2.5 <sup>c</sup>                |
| Metabolic myopathies             | 0.8                             |
| Myasthenia gravis                | 0.5                             |

<sup>a</sup>Prevalence figures related to male population, expected to represent 50% of the total population.

<sup>b</sup>The one female symptomatic carrier is not included in the number.

<sup>c</sup>Limb girdle type 2 I made up 60% of the total number.