## The incidence and distribution characteristics of MLL rearrangements in Chinese acute myeloid leukemia patients by multiplex nested RT-PCR

Hua Yang<sup>a,\*</sup>, Tingting Cao<sup>a</sup>, Li Gao<sup>b</sup>, Lili Wang<sup>a</sup>, Chengying Zhu<sup>a</sup>, Yuanyuan Xu<sup>a</sup>, Yu Jing<sup>a</sup>, Haiyan Zhu<sup>a</sup>, Na lv<sup>a</sup> and Li Yu<sup>a,\*</sup>

<sup>a</sup>Department of Hematology, The Chinese PLA General Hospital, Beijing 100853, China <sup>b</sup>Department of Hematology, China-Japan Friendship Hospital, Beijing 100029, China

This article has been retracted. You can find the retraction notice here: https://content.iospress.com/ articles/technology-and-health-care/thc209060.

DOI: 10.3233/THC-171329

Journal: Technology and Health Care, vol. 25, no. S1, pp. 259-265, 2017

Published 20 July 2017

0928-7329/17/\$35.00 (c) 2017 - IOS Press and the authors. All rights reserved

This article is published online with Open Access and distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC-BY-NC 4.0).

<sup>\*</sup>Corresponding authors: Hua Yang, Department of Hematology, The Chinese PLA General Hospital, 28 Fuxing Road, Beijing 100853, China. Tel.: +86 13260373275; E-mail: yananshen2006@163.com; Li Yu, Tel.: +86 010 55499003; Fax: +86 010 68150721; E-mail: chunhuiliya@yahoo.com.